

Micro-Raman spectroscopy of the shocked olivine in the Martian meteorite ALH 77005

Szabolcs Nagy (1) (ringwoodit@yahoo.com), Sándor Józsa (1), Arnold Gucsik (2), Szaniszló Bérczi (1), Kiyotaka Ninagawa (3), Hirotsugu Nishido (4), Miklós Veres (5), Ákos Kereszturi (1), Henrik Hargitai (1)

- (1) Eötvös Loránd University of Budapest, H-1117 Budapest, Pázmány Péter sétány 1/c, Hungary
- (2) Max Planck Institute for Chemistry, Department of Geochemistry, Becherweg 27, D-55128, Mainz, Germany
- (3) Department of Applied Physics, Okayama University of Science, 1-1 Ridai-cho, Okayama, 700-0005, Japan
- (4) Research Institute of Natural Sciences, Okayama University of Science, 1-1 Ridai-cho, Okayama, 700-0005, Japan
- (5) Research Institute of Solid State Physics and Optics of the Hungarian Academy of Sciences, H-1525 Budapest, Konkoly Thege Miklós út 43-49, Hungary

Introduction:

Olivine is one of the most refractory and abundant igneous rock-forming minerals, which has been proven useful as an indicator of shock metamorphism in the study of meteorites [1] [2]. The systematic studies of micro-Raman properties of experimentally shocked olivine crystal assemblages have been performed by [3] and [4]. [3] have found no significant characteristic relation between Raman shifts as a function of increasing shock pressure in olivine. However, [4] described a new metastable phase formed at ambient pressure as the metastable intermediate during back-transformation from wadsleyite.

Sample and methods:

Two thin sections (ALH 77005-A and ALH 77005-B) were supplied from National Institute of Polar Research (NIPR, Tokyo, Japan), which belong to the meteorite collections. They were prepared as polished thin sections.

The ALH 77005 Martian meteorite is a lherzolithic type shergottites. Olivine (Fo72) occurs as an anhedral to subhedral grains up to 2 mm in length. The ALH 77005 consists of melt pockets, which are mostly crystallized as the spinifex texture. Olivine in the bulk of ALH 77005 has brownish color because of approximately 4.5 wt% of the total iron (Fig. 1).

Raman spectra were recorded with a Renishaw Rm-2000 Raman spectrometer attached to a Leica DM/LM microscope. The 785 nm line of the diode laser served as excitation source, the laser power on the sample was 8 mW. The microscope has focused the excitation beam into a 1 μm diameter spot. Raman spectra were recorded in the 200-1100 cm^{-1} region using a CCD camera, the accumulations lasted during 60 sec.

Results and discussion:

We have investigated the olivine grains based on the increasing distance from the melt pocket to the outermost position of the sample. Close to the melt

pocket (sp1), the Raman spectrum shows not significant changes (Fig. 2). In the intermediate region (sp2) a new peak appears at 755 cm^{-1} , and observed a slightly broadening features (Fig. 2).




Fig. 1 Brown olivine grains in the ALH 77005.

In the third spectrum (sp3) the 416 cm^{-1} peak disappears (Fig. 2). The disappearance of the olivine-related peak at 600 cm^{-1} may indicate the presence of a new phase such as $\xi\text{-(Mg, Fe)}_2\text{SiO}_4$ [4]. In this spectrum a medium peak at 535 cm^{-1} is presented, which may be related to the $\xi\text{-(Mg, Fe)}_2\text{SiO}_4$ phase. This peak is not related neither to α and β nor γ -olivine phases. In this spectrum there is an appearing peak at 918 cm^{-1} , which is related to the wadsleyite. In the fourth spectrum (sp4) which is lie the farthest point from the melt pocket, the 600 cm^{-1} olivine peak is absolutely disappearing.




Fig. 2 Raman-spectra of different distance of olivine grains from the melt pocket.

Conclusion:

These Raman-spectra indicate a belt-like arrangement of the structural changes of olivine grains from the melt pocket away to the far regions of the sample.

Acknowledgement:

Authors are grateful to Dr. H. Kojima for the loan of the Antarctic Meteorite set (NIPR, Tokyo, Japan).

This study has been supported by the international joint collaboration of JSPS-HAS (No. 2007/104).

References:

- [1] Stöffler, D. (1974) Deformation and transformation of rock-forming minerals by natural and experimental processes: II. Physical properties of shocked minerals. *Fortschritte der Mineralogie*, 51, pp 256-289.
- [2] Boctor, N. Z., Fei, Y., Bertka, C. M., Alexander, C.M.O.D., and Hauri, E. (1999) Shock metamorphic effects in Martian Meteorite ALH 77005 XXX LPSC, #1628.
- [3] Farrell-Turner, S., Reimold, W. U., Nieuwoudt, U. and Erasmus, R. M. (2005) Raman spectroscopy of olivine in dunite experimentally shocked to pressures between 5 and 59 GPa. *Meteoritics & Planetary Science*, 40, pp 1311-1327.
- [4] Van de Moortéle, B., Reynard, B., McMillan, P. F., Wilson, M., Beck, P., Gillet, P., Jahn, S. (2007) Shock induced transformation of olivine to a new metastable $(\text{Mg}, \text{Fe})_2\text{SiO}_4$ polymorph in Martian meteorites. *Earth and Planetary Science*, 261, pp 469-475.